

The client sample was analyzed for plant-based cannabinoids by Liquid Chromatography (LC). The collected data was compared to data collected for certified reference standards at known concentrations.

113/20-01			
ID	Weight %	Concentration (mg/g)	
<b>Δ9-THC</b>	0.0776	0.776	
THCV	ND	ND	
CBD	1.35	13.5	
CBDV	ND	ND	
CBG	ND	ND	
CBC	0.0588	0.588	
CBN	ND	ND	
THCA	0.518	5.18	
CBDA	17.4	174	
CBGA	0.608	6.08	
CBDVA	0.107	1.07	
∆8-THC	ND	ND	
exo-THC	ND	ND	
Total	20.1	201	0% Cannabinoids (wt%) 17.4%
Max THC	0.532	5.32	Limit of Quantitation (LOQ) = 0.0067 wt%
Max CBD	16.6	166	Limit of Detection (LOD) = $0.0022 \text{ wt\%}$

## Ratio of Total CBD to THC 31.2:1

Max THC (and Max CBD) are calculated values for total cannabinoids after heating, assuming complete decarboxylation of the acid to the neutral form. It is calculated based on the weight loss of the acid group during decarboxylation: MAX THC =  $(0.877 \times THCA) + THC$ . This calculation does not include other cannabinoid isomers (eg. D8-THC and exo-THC). ND=None detected above the limits of detection (LOD), which is one third of Limit of Quantification (LOQ). For values reported as "<LOQ", the estimated value is included in the calculated Total.

## **END OF REPORT**

113720-CN